

LOVES TO EAT THIS CEREAL, IT CHECKED HER CONSTIPATION*

Kellogg's ALL-BRAN Helped Miss Kesterke

We quote from her letter: "Three years ago, I became constipated. I tried many laxatives. But as soon as I got used to each kind, I began to get the same trouble. Last summer I was on my vacation. They served Kellogg's ALL-BRAN. I just loved it. I eat Kellogg's ALL-BRAN every morning, and ever since I have not had to take any more laxatives."

*Due to insufficient "bulk" in meals. Tests show Kellogg's ALL-BRAN provides "bulk" to aid elimination. ALL-BRAN is also rich in vitamin B and iron.

The "bulk" in ALL-BRAN is gentle. It resists digestion better than the fiber in fruits and vegetables, so it is often more effective.

Isn't this food safer than risking patent medicines? Two tablespoons of ALL-BRAN daily are usually sufficient. If seriously constipated, use with each meal. See your doctor, if you do not get relief.

Use as a cereal, or in cooking. Sold by all grocers. Made by Kellogg in London, Ontario.

Keep on the Sunny Side of Life

HOLSTEIN ASSN. HOLD ANNUAL FIELD DAY

A large number of Holstein breeders and others interested in dairying attended the annual field day of the Holstein Friesian Association at Mr. R. A. Profit's, Freetown, on Saturday. The afternoon's program included cattle judging competitions by boys and girls and short addresses by the President of the Association, Mr. Colby Lewis, Freetown; Mr. Allison Profit, Freetown; Mr. Cecil Stewart, Hampshire; Mr. W. R. Shaw, Deputy Minister of Agriculture, Cornwall; Mr. W. L. Brenton, Dairy Superintendent, Charlottetown; Louis Cairns, Freetown; Mr. J. W. Nicholson, Bedouque; Mr. A. McLean, M.P.; Dr. J. A. Clark, Superintendent of the Experimental Station, Charlottetown, and Mr. Louis Roper of the Dominion Live Stock Branch, Charlottetown.

Sport competitions followed the regular field day program. The prize winners in the cow judging competition were as follows: 1, Clifton Stewart, Hampshire; 2, Henry Godfrey, North Wilshire; 3, Cecily Leard, Tryon; 4, Arnold Godfrey, North Wilshire; 5, Alice Fell, Tryon. In the heifer judging competition the winners were: 1, Henry Godfrey, North Wilshire; 2, Cecily Leard, Tryon; 3, Arnold Godfrey, North Wilshire; 4, Jessie Leard, Tryon; 5, Albert MacNutt, Malpeque.

The winners in the various sport events were as follows: 50 yard dash for girls under 9: 1, Eleanor Harvey, Carleton R. R.; 2, Laura Hill, Freetown; 3, Marion Leard, Tryon. 50 yard dash for boys 9-12: 1, Boyd Beardo, Freetown; 2, Buddy Shaw, Cornwall; 3, Robert Cairns, Freetown. 50 yard dash for girls 12-15: 1, Jessie Leard, Tryon; 2, Alice Fell, Tryon. 100 yard dash for boys 12-15: 1, Claude Lewis, Freetown; 2, Clifton Stewart, Hampshire; 3, Boyd Beardo, Freetown. 100 yard dash for boys over 15: 1, Allan Clow, Freetown; 2, Walker Stewart, Freetown; 3, Ralph Harvey, Carleton. Ladies free for all, 100 yards: 1, Dorothy Mayne, Springfield; 2, Jessie Leard, Tryon; 3, Mae Gardner, Freetown. Men's free for all, 100 yards: 1, Arnold Clow, Freetown; 2, Albert MacNutt, Malpeque; 3, Ina Lewis, Freetown. 25 yard dash for boys under 9: 1, Bobbie Reid, Kinkora; 2, Edwin Lewis, Freetown; 3, Earl Stewart, Hampshire.

Mr. Colby Lewis, Freetown, President of the Association, in a short address told something of the work of the organization. Of 500 cattle tested for Bangs Disease last year only 4 reacted. The diseased cattle have since been slaughtered. He urged that more breeders join up with the Association in order that more and better work might be done. An address of welcome to the members of the Holstein Friesian Association and all others present was delivered by Mr. Allison Profit.

Mr. Cecil Stewart, Secretary of the Association, reported that Mr. W. R. Shaw, Deputy Minister of Agriculture, in a short talk referred to the many problems facing the farmer. The biggest problem facing the cattle breeders is that of nutrition. A few years ago the idea prevailed that the farmer should import feeds in large quantities in order to make a success of dairying. There was very little attempt to investigate the possibility of local feeds. One thing, perhaps, that the depression has done has been the impetus it has given to the investigation of the possibilities of the home grown feeds such as alfalfa, pea, clover and vetch mixtures, corn, etc. It has been found that there is very little need to spend money on imported concentrates. The Department of Agriculture has been giving assistance in the investigation of the value of home grown feeds as well as in many other matters.

The speaker congratulated the Holstein Friesian Association for the assistance it gave in the promotion of the tuberculosis tests some years ago. He expressed the hope that at some time in the future a general test might be made for Bangs Disease.

He referred to the warble fly as a pest in the control of which all dairymen should be vitally interested.

Mr. W. L. Brenton, Dairy Superintendent, spoke briefly. What is needed in the dairy industry in this Province is not so much an increase in the number of units of production as an increase in the production of each unit. He emphasized the need for great care in the handling of milk and cream. The quality of butter and cheese depends to a large extent on the quality of the raw material, he said.

Dr. J. A. Clark spoke on pasture improvement. He urged the dairymen to pay greater attention to pastures as one means of increasing summer production of milk. A vote of thanks to Mr. and Mrs. Allison Profit and Mr. Profit, Sr., for their generosity in allowing the Association the use of their premises was moved by Mr. Cecil Stewart, seconded by Mr. Granville Buntain.

WILL SPEND MONTH ON P. E. ISLAND (C.P. By Guardian's Special Wire) SYDNEY, July 7—Rev. E. M. Aitken, B. A., pastor of St. Andrew's United Church here, will leave early this week for Prince Edward Island where he will remain for about a month. From there he will proceed to Calgary to take up his new duties as pastor of Central United Church. Rev. Mr. Aitken's resignation from St. Andrew's Church becomes effective on July 31st.

Trinity United Church Hogg - Cousins Wedding

MONDAY, JULY 8 7:30 p.m.—Committee of Stewards, Board Room.

The Central Guardian

This column is reserved for news of local interest but advertising of a new nature may be inserted at a cents a word strictly payable in advance.

W. K. ROGERS Agencies Limited, general insurance agents are now to be found in their new quarters in the Currie Building at 181 Queen Street. L-8506-7-3-61.

THERE WILL BE a meeting of the First Aid Class at the Drill Hall at 7 o'clock tonight. Tomorrow night at the same time examinations for certificates will be held. L-3672.

VISITING HERE—Mrs. Thomas Keoughan of 52 Weymouth St., arrived home from Boston where she attended the ordination of her nephew Rev. James McWade, grandson of the late James J. McWade of St. Teresa. She was accompanied by Mrs. Michael J. McWade and daughter Dorothy, mother and sister of Rev. Fr. McWade. They will be the guests of Mr. and Mrs. Keoughan for the summer.

UNUSUAL OPERATION—An unusual operation was performed recently in the Prince Edward Island Hospital to save the life of the infant daughter of Mr. and Mrs. Carl Cameron, City. The operation was made necessary by a rare condition, the growth of a tumor in the intestines, and was performed successfully by Dr. Donald Campbell. Up till July last year medical journals recorded only 64 such operations.

PERSONALS Mr. Kenneth Leslie, Halifax, is visiting Charlottetown, the guest of Lt.-Col. and Mrs. K. S. Rogers.

Mr. and Mrs. J. M. Murley leave this morning on a short visit to Halifax.

Prof. A. R. Weir of Saskatoon University, with Mrs. Weir and family have arrived by motor from Saskatchewan to spend the summer with Mrs. Weir's parents, Mr. and Mrs. G. H. Taylor, Charlottetown.

Mr. George H. Scott, Acting Provincial Commissioner of the Boy Scouts of New Brunswick, and Mr. Joseph Guss, Brunswick Secretary, attended the various functions held in honour of Lord and Lady Baden-Powell in Charlottetown last week.

Mr. A. D. Merkel, Atlantic Superintendent, Canadian Press, Halifax, accompanied by Mrs. Merkel, their daughter, Miss Peggy and Miss Mary Elizabeth Merkel, at Miss Helen Connor, Halifax, arrived Saturday to spend a vacation at Holland Cove.

Misses Audrey MacKay, Evelyn Harvey, Jean Bonifer, Helen Keough, Marjory Donnelly, Mary Tierney and Mr. Sutherland MacKay, all of Albany, were the guests of Miss Audrey Harvey, Linkletter, last week.

REGINA Free Of Camp Strikers (C. P. by Guardian's Special Wire) REGINA, July 7—Regina, free of the army of relief camp strikers that made the city their headquarters for three weeks, breathed easier today as members of the scattered battalions made their way homeward. The move left Winnipeg the only Western city in which relief camp strikers are gathered in any appreciable number.

There the men remained firm in their intention of presenting their complaints against camp conditions to Ottawa authorities. Some 600 of them camped under canvas in the exhibition grounds and were being fed by the Manitoba Government.

Approximately 125 men detained at Calgary to be met by a small crowd. Everything was orderly. The men were jolly and in good spirits.

Seventy-six of those at Calgary however, refused to register for camp work and officials announced no further aid would be extended. Others went to their homes or to friends.

And, in the city they left, a Saskatchewan Government commission prepared to launch an investigation into the causes and circumstances of the fatal Dominion Day riot.

Detective Charles Millar was killed, and 100 injured in the fighting between strikers, citizens and police. Many were arrested, including Arthur Evans, strike leader.

Sixty of those taken into custody were freed Saturday and seven others were remanded when they appeared in court. It was estimated 61 others remained in custody. The condition of those injured in the riots and still in hospital, was reported improving.

Some 150 strikers whose homes were east of Regina arrived in special railway coaches at Winnipeg where 24 of them detoured east to Ontario, Quebec and the Maritimes.

Offer of 600 jobs on farms and camps were given Manitoba's quota of camp strikers Saturday but so far none has taken advantage of them.

—EXCURSION— PICTOU CARNIVAL, JULY 10TH Boat leaves Bruce Stewart's Wharf at 7 a.m. Get your tickets at Hughes' or Old Spain. L-8667-7-8-21

THE FLAVOR LASTS



it's always fresh - it keeps you fresh

RELIGIOUS SERVICES

EVANGELISTIC LECTURE Handled. At the evening service the members of the Royal Orange Association attended in a body and occupied the front pews in the centre section. They were welcomed by Rev. Mr. Miller who preached from the text Mark 12:17, "And Jesus answering said unto them, render to Caesar the things that are Caesar's and to God the things that are God's, and they marvelled at Him." It was a most appropriate sermon and eloquently delivered. A large congregation was present and the visiting brethren made a most creditable appearance.

After the opening prayer Trinity's double quartette sang impressively the 23rd Psalm.

HOLY REDEEMER CHURCH

Very practical sermons were preached by Rev. W. E. Enright, C.S.S.R., at all the Masses yesterday morning. In his discourse, Father Enright lucidly explained the necessity of vacations for both children and adults, and exhorted the parishioners to spend their vacations in a Christian manner. Persons who ignore God during their vacation and resume pious practices after their vacation is over are insincere in their piety, Father Enright emphasized. He also reminded the parents of their duties towards their children during the vacation season, and urged them to guard closely the spiritual welfare of the children during the holidays.

Masses were at the usual Sunday hours, High Mass being celebrated by Rev. C. O'Hara, C.S.S.R.

Devotions in honor of Our Mother of Perpetual Help, which are held on the first Sunday of each month, took place last evening.

CONSPIRACY

(Continued from Page 1) from Sydney to Digby. In January 1933 dorries from the Nashwaak went ashore at Glace Bay with kegs that had been picked up at sea, Burns said. In March of the same year dorries contacted the Nashwaak and made landings at Bear River, N. S.

"You were paid always by Captain Mosher," said W. G. Ernst, M.P., one of the defence lawyers.

"Yes," "Mr. Ritcey was never present?" "No."

Copies of telegrams purporting to tell of dealings between several of the accused and liquor agents were shown by J. H. Trappell, manager of the Canadian National Telegraphs in Halifax, who appeared on a subpoena.

REQUESTS TO POLICE HOVE, England—In appreciation of their protection Mrs. N. T. Welch a widow, bequeathed \$500 each to two policemen whose beats passed her home.

WOMEN THREATEN FAST LYALLPUR, India—Forty Sikh women have made a pact to fast to death unless certain warring factions of the community do not come to an understanding.

TRIED TO BURN EVIDENCE LONDON—Two years imprisonment was imposed on Fred Wood a cashier who pleaded guilty to concealing fire to premises to conceal evidence of a deficiency in his accounts.

SIGNING UP FOR PEACE LONDON—Nine million people have already signed an "annual peace ballot" being circulated by peace societies throughout the United Kingdom.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.

Control of Black Rot of Apples

(Experimental Farms Note) Black rot is proving to be an important disease of apples in Eastern Canada. It is caused by a fungus which is capable of attacking the trunk, limbs, leaves and fruit. When the trunk and limbs are involved, located in the following manner: soon after they unfold from the buds. The first symptoms appearing on the foliage are purplish spots which gradually become brown with a light grey centre and faintly raised margins. Owing to the fact that the fungus usually gains entrance through wounds, caused by insects or other mechanical injuries, there has been much winter injury in many orchards during the last two years, trees so affected are more susceptible to attack. Observations made by the Dominion Field Laboratory of Plant Pathology at Fredericton, New Brunswick, show that the following measures should be kept in mind as a basis for prevention of the disease. Close pruning to remove all cankers and dead wood and clean cultivation or ploughing under all dead leaves are commendable practices. All prunings diseased bark and mummies to be destroyed. The ground should be destroyed, preferably by burning, because they serve as sources from which the disease may spread. Pruning wounds or other mechanical injuries that cannot be avoided should be treated with a good disinfectant, such as Bordeaux mixture. While there is no positive evidence that standard fungicides such as Bordeaux mixture and lime sulphur, satisfactorily control the disease, it is nevertheless advisable to keep the foliage and limbs constantly covered with spray to secure at least a partial protection. Attention should be given to the control of insect pests and other fungous and bacterial diseases which result in wounds that may serve as avenues for infection. Further information on this disease can be obtained from nearest Plant Pathological Laboratory.